Caution: SabrEx for Soybeans In-Furrow may be used on either untreated seed or seed that has been treated with a fungicide or insecticide. For seed treated with a fungicide or insecticide, wear appropriate PPE to avoid direct skin contact, and follow the caution statements of the other treatments. In case of direct contact with skin or eyes, immediately flush the exposed area with water. Seek medical attention if irritation develops or persists.

Warranty: Advanced Biological Marketing guarantees the cultures in this product will inoculate the quantity and kind of seed specified on this package when: 1) Product is used before the expiration date; 2) Product is stored properly; 3) Product is used under favorable soil and weather conditions; 4) Product is used in accordance with directions for use noted on this package and other technical recommendations referenced on this label. All field performance complaints must be brought to the attention of Advanced Biological Marketing within seventy (70) days of planting. ADVANCED BIOLOGICAL MARKETING MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS, WARRANTY OF MERCHANTABILITY, OR WARRANTY OF A PARTICULAR PURPOSE, OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. ADVANCED BIOLOGICAL MARKETING'S LIABILITY IS HEREBY EXPRESSLY LIMITED TO THE PURCHASE PRICE OF THIS PACKAGE ONLY. IN NO EVENT SHALL ADVANCED BIOLOGICAL MARKETING OR ANY SELLER OF THIS PRODUCT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.


SabrEx for Soybeans In-Furrow is intended to utilize Trichoderma's ability to improve nutrient availability to plants.

Contains living organisms. Store in cool dry place out of direct sunlight 40°F to 77°F. Do not freeze.





Lot number:





Application Rate: 1 fl. oz. per acre. This jug treats 128 acres. Rate per 1000 row feet is dependent on row spacing.

Directions for Use:

Apply at planting time as an in-furrow application by directing into the furrow before the seed is covered.

Mixing Instructions:

Agitate a thoroughly clean tank half full of carrier. Maintain constant agitation. Add SabrEx for Soybeans In-Furrow liquid. Add remaining carrier.

SabrEx for Soybeans In-Furrow can be tank mixed with most recommended in-furrow products. A compatibility or jar test is recommended when using anything besides water as the carrier.

Tank Mix Restriction:

SabrEx for Soybeans In-Furrow can be left in the tank mix for a maximum of 24 hours; longer may result in loss of viability. IF THIS TIME IS EXCEEDED, DISCARD THE MATERIAL OR ADD MORE SABREX FOR SOYBEANS IN-FURROW.

CONTAINS NON-PLANT FOOD INGREDIENTS

Guaranteed Analysis:

Active Ingredients: 0.10% - *Trichoderma virens* 1 x 10⁸ cfu/ml Inert Ingredients: 79.9%-Water, 20.0%-Aqueous solution