Contents settle - shake well before measurement and use.





GH2 SIGNAL WORD: Danger

SKIN: Wash with plenty of soap and water. IF INHALED: Remove person to fresh air and keep or in a well-ventilated area. Wear protective gloves, respiratory protection, eye protection. IF ON breathing dust. Wash face, hands, and any exposed skin thoroughly after handling. Use only outdoors

local, regional, national, and international regulations. contaminated clothing and wash it before reuse. Dispose of contents/container in accordance with

DIRECTIONS FOR USE: This product is intended for nitrogen fixation and to improve nutrient

availability. Graph-Ex SA for Soybeans may be applied to seed using two methods of application,

BULK APPLICATION: Apply one half (0.5) ounce by weight per 140k unit of seed, by manually or

mechanically dispersing Graph-Ex SA for Soybeans into a flowing stream of seed. Ensure the product is

evenly dispersed in the planter or drill box.

140k unit increments.

POSSIBLE, RE-INOCULATE BEFORE PLANTING.

information, please contact your Graph-Ex SA representative at 877-617-2461. For current "planting window" recommendations, and the latest seed treatment compatibility

recommendations referenced on this label. All field performance complaints must be brought to the attention of Agrauxine Corp. within seventy (70) days of planting. AGRAUXINE CORP. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITUESS, WARRANTY OF MERCHANTABILITY, OR WARRANTY OF FITUESS, WARRANTY OF FITUESS, WARRANTY OF FITUESS, WARRANTY OF FITUESS. conditions; (4) Product is used in accordance with directions for use noted on this package and other technical and inoculants. Topsoil exceeds 80°F • Soil erosion • Use of soil treatments and chemicals injurious to soil bacteria

LIMITED WARRANTY: Agrauxine Corp. guarantees the culture in this product will inoculate the quantity and kind of seed specified on this package when: (1) Product is used before the expiration date; (2) Product is stored properly as prescribed by the label; (3) Product is used under normal farming practices with favorable soil and weather (1) product is used under normal farming practices with favorable soil and weather as prescribed by the label; (3) Product is used under normal farming practices with favorable soil and weather

FROM THE USE OR HANDLING OF THIS PRODUCT. SELLER OF THIS PRODUCT BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FIMILED TO THE PURCHASE PRICE OF THIS PACKAGE ONLY, IN NO EVENT SHALL AGRAUXINE CORP, OR ANY PURPOSE, OR ANY COMPANY'S LIABILITY IS HEREBY EXPRESSLY

Soil Amending Guaranteed Analysis Active Ingredients:

1.2%

UNIAINS NUN-PLANT FUUD INGREDIEN

Shake well before measurement and use.

<mark>Induc</mark>ed Gene Expression Triggers

Net Weight: 3 oz. (85 grams) Treats 6 x 140k units of seed

Contents settle - shake well before measurement and use.

Agrauxine by Lesaffre

AGRAUXINE CORP. P.O. Box 222 | Van Wert, OH 45891

STORAGE INSTRUCTIONS: Contains living organisms. Store packaged in a cool dry place between 40° and 77°F and keep closed until use. Use inoculant before expiration date and only use on specified legume seed. Protect inoculant and inoculated seed from direct sunlight. Information regarding the contents and levels of metals in this product is available on the internet at aapfco.org/metals

Use before expiration date printed on this container. EXP. DATE:

Graph-Ex Inoculant Seed Lubricant

for **Soybeans**

with

.....Microbial Confent Bradyrhizobium Japonicum 7.5 x 10° cfu/g Trichoderma virens 2.0 x 10° cfu/g

Total Inert Ingredients (lubricant, carrier, stabilizers)

.%8.8%

LOT NUMBER:

 fould play a proper than 8.5
Oreganic matter is less than 4.5.
Drought or fload a proper than 8.5. than three (3) years • Soil pH is less than 5.8 (the pH should be adjusted by liming prior to inoculation) second inoculant to achieve at least a double rate. • Soil has not hosted the specific legume for more combination of inoculants. American Best Inoculant Sterile Peat is highly recommended to be used as a any of these conditions are present, Agrauxine Corp. highly recommends at least a double (2x) rate of a KECOMMENDATIONS FOR ADVERSE CONDITIONS OR NEW "VIRGIN" SOYBEAN FIELDS: When

into planter or drill box. Mix seed until uniform coverage is obtained. It is best to treat and mix seed in PLANTER BOX APPLICATION METHOD: Apply one half (0.5) ounce by weight per 14-0k unit of seed

bulk or planter box.

PRECAUTIONARY STATEMENTS: Obtain, read, and follow all safety instructions before use. Avoid

inhalation. Causes skin and eye irritation. May cause respiratory irritation. HAZARD STATEMENTS: May form combustible dust concentrations in air. May cause cancer through

FOR OPTIMAL PERFORMANCE, PLANT SEED WITHIN 72 HOURS OF APPLICATION. IF NOT

comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Gently remove contact lenses, if present. Continue rinsing. If unwell or concerned: Get medical advice. Take off