



Soybean Inoculant Pesticide Compatibility, Planting Window and Tank Mix Recommendations

ABI PLUS 30 Extender PLUS America's Best Soybean Inoculant PLUS pesticide: Maximize Planting Window Allowed After Inoculant Application							
Liquid Soybean Inoculant Formulations with ABI Plus 30 Extender		Pesticide(s)	In Tank Slurry ¹	Maximizing Planting Window (Days)			
				1 Day	7 Days	21 Days	28 Days
		No Pesticide	24 hours	28 Days			
 ABI™ Sterile Liquid 10L Soybean ABI™ Sterile Liquid 3.1L Soybean MegaPack™ Low Volume Liquid	with ABI Plus 30 Extender	ApronMaxx® RFC ApronMaxx® RTA® Innovate® System Warden™ RTA®	4 hours	28 Days			
		Acceleron® ApronMaxx® BeanPak™ CruiserMaxx® CruiserMaxx® Advanced Pioneer PPST Poncho® / VOTIVO® Trilex® 6000	4 hours	21 Days			
		Any Treatment Containing Molybdenum	DO NOT TANK MIX	1 day			

America's Best Inoculant PLUS pesticide: Maximize Planting Window Allowed After Inoculant Application									
Liquid Soybean Inoculant Formulations WITHOUT EXTENDER		Pesticides (s)	In Tank Slurry ¹	Maximizing Planting Window (Days)					
				1 Day	2 Days	3 Days	4 Days	5 Days	6 Days
		No Pesticide	24 hours		7 Days				
 ABI™ Sterile Liquid 10L Soybean ABI™ Sterile Liquid 3.1L Soybean MegaPack™ Low Volume Liquid		ApronMaxx® RFC ApronMaxx® RTA® CruiserMaxx® CruiserMaxx® Advanced Pioneer PPST Poncho® / VOTIVO® Innovate® System Warden™ RTA®	4 hours	5 Days					
		Acceleron® ApronMaxx® BeanPak™ Trilex® 6000	4 hours	3 Days					
		Any Treatment Containing Molybdenum	DO NOT TANK MIX	1 day					

Soybean Inoculant Pesticide Compatibility, Planting Window and Tank Mix Recommendations

America's Best Inoculant PLUS Pesticides: Maximum Planting Window Allowed After Inoculant Application		
Dry Soybean Inoculant Formulations	Inoculant Only	Inoculant plus one of: Acceleron[®], ApronMaxx[®] RFC, ApronMaxx[®] RTA[®], CruiserMaxx[®], CruiserMaxx[®] Advanced, Poncho[®] / VOTiVO[®], Innovate[®] System, Pioneer PPST, Warden[™] RTA[®]
ABI Non-Sterile Peat Soybean	Plant within 24 hours of application	Plant within 4 hours of application
ABI Sterile Peat Soybean	Plant within 24 hours of application	Plant within 24 hours of application

Precautions: Store treated seed in cool place, out of direct sunlight. Always use non-chlorinated water for pesticide dilutants when mixing with inoculants. Read and follow all label directions.

"Slurry"¹ refers to co-mixing liquid inoculant together with the seed treatment in the treater tank. If possible avoid co-mixing liquid inoculant together with the pesticide seed treatment, use a separate tank for inoculant. The inoculant plus the named seed treatment should be kept together in the tank no longer than four hours before application to the seed.

America's Best Inoculant[®] for Soybeans is a registered trademark of Advanced Biological Marketing. Acceleron[®], Apron-Maxx[®] RFC, ApronMaxx RTA , ApronMaxx[®] Bean Pak, CruiserMaxx[®] are trademarks or registered trademarks of Syngenta Group Company. Warden[™] RTA[®] is a trademark of Agrilience LLC. Poncho[®]/VOTiVO[®] and Trilex[®] are registered trademarks of Bayer Crop Science.